



## A prospective study of patients with Blunt Abdominal Trauma: A study from the northern part of India

Dr. Sandesh Kumar Shrivastava<sup>1</sup>, Dr. UC Singh<sup>2\*</sup>

<sup>1-2</sup> Professor, Department of Surgery, BRD Medical College, Gorakhpur, Uttar Pradesh, India

\*Corresponding Author: Dr. UC Singh

### Abstract

**Background:** Blunt abdominal trauma (BAT) has become a major health problem all around the world mainly in low and middle-income countries.

**Aims and Objective:** To study the incidence, reasons and outcome of BAT in order to diagnose it properly and to start early intervention to decrease associated morbidity and mortality.

**Materials and Methods:** It was a prospective study including patients who presented with history of blunt trauma to abdomen in the Department of Surgery, BRD Medical College Gorakhpur from January 2012 to August 2014. Detailed history, mode of trauma, treatment given, extra abdominal injuries, clinical presentation, internal organ injuries, hospital stay, associated complications and morbidity parameters were studied in all the patients.

**Results:** A total of 134 patients were admitted during the study period. Incidence of BAT was 54/ 6204, 50/6794 and 30/5572 in 2012, 2013 and 2014 respectively with male predominance involving mostly patients with age group between 20-30 years. Road traffic accidents were the most common cause of BAT. Most of the patients in 2012 were treated conservatively but in 2013 and 2014 surgical intervention were most common. Injuries of Head and chest were the common extra abdominal injuries, in abdomen small intestine was damaged in majority of patients. Most of the patients stayed for 7 to 12 days in the hospital and though 24 (17.91%) patients died due to severity of the trauma, 104 (77.61%) recovered without any complications.

**Conclusion:** The BAT is preventable as the road accidents are the major cause. Sound judgment with effective treatment and surgical expertise can improve the survival of patients with BAT.

**Keywords:** blunt abdominal trauma, road traffic accidents, abdominal injuries

### Introduction

Trauma especially road traffic injury is one of the major health related issue throughout the world especially in developing countries like India <sup>[1]</sup>. Road traffic accidents still remains the one of the top reason for blunt abdominal trauma (BAT), other causes of BAT include recreational accidents, fall and assault <sup>[2]</sup>.

BAT is very common and the prevalence of intra-abdominal injury after BAT range from 12% to 15% <sup>[3, 4]</sup>. Diagnosis of BAT can be done using physical examination, ultrasonography, computed tomography (CT), laparoscopy, laparotomy and using different laboratory tests <sup>[5]</sup>.

Data from the different study had shown a high mortality in patients with BAT especially when patients is having multiple solid organ injury <sup>[6, 7]</sup>. During surgery risk of death is increases by 4.4 times in patients with solid organ injury <sup>[7]</sup>. In order to salvage patients from such abdominal trauma, rapid resuscitation is needed <sup>[6]</sup>.

Several old studies have recommended an observation period of 23 hours following BAT; nevertheless the optimal observation duration is still unknown <sup>[8]</sup>.

Hence, present study was done to find out the incidence, reasons and outcome of BAT in order to diagnose it properly and to start early intervention to decrease associated morbidity and mortality.

### Materials and Methods

The present prospective study included patients who presented with history of blunt trauma to the abdomen, in the Department of Surgery, Department of Surgery, BRD Medical College Gorakhpur from January 2012 to August 2014.

A Written informed consent from all the patients and Ethical Committee approval was obtained before starting the study.

Detailed history of patients including the length of delay in treatment taken and nature of trauma including weapon responsible for trauma was recorded. Conscious patients without any alcoholic and narcotic drug consumption history were asked the detailed history including any presenting symptoms and exact site of injury.

Those patients who sustained injury due to assault by fists and blows, lathi, iron rods/kicks or received accidental animal kicks over abdomen or got injured due to fall of some heavy object like log, gravel, got buried in mine or had rolled down the stairs or stumbled over some blunt object and were resuscitated in the casualty and admitted to the wards and subjected to appropriate management and the admitted patient who died during the treatment were included in the present study.

Patient who reported to hospital with history of blunt abdominal trauma but on examination showed no distant symptoms and sign of abdominal injury, and/or given

conservative line of treatment, and in due course showed improvement, without any deterioration and not admitted in hospital were excluded from the present study.

If patient was in the state of shock often with presence of internal injury, his cardio respiratory status was assessed and recording of vital sign was made immediately and in order of priority, establishment of adequate ventilation, control of major hemorrhage done.

Insertion of central line and cross match for blood transfusion was done simultaneously. Rapid I.V. bolus of crystalloid solution, dextran and blood was given as required.

With respiratory and circulatory efficiency resorted, vital sign recording continued at 15 minute intervals. Detailed physical examination and local examination was performed.

The wound was covered by occlusive dressing. Foley's catheter inserted for recording hourly output and urine sampling to evaluate the integrity of urinary track. A Ryle's tube inserted for nasogastric suction whenever necessary in abdominal injury.

Resuscitation was continued throughout the period of evaluation until a treatment plan, surgery or observation was established.

The external injury was noted in detail and finding such as surgical emphysema, active bleed peritoneal/pleural breach, evisceration, guarding & rigidity in abdomen on palpation were recorded.

Laparotomy was performed on selective patients who showed parietal peritoneal breach, evisceration of either bowel or omentum and any foreign body felt per abdomen.

All the data were analyzed using IBM SPSS- ver.20 software. Analysis was performed using chi-square test and independent sample student t test. P values <0.05 was considered to be significant.

## Results

In present study out of 6204, 6794 and 5572 patients admitted 54, 50 and 30 patients were found to have BAT (incidence being 0.87%, 0.74% and 0.54% respectively) in 2012, 2013 and 2014 respectively.

Out of 134 BAT patients, 112 (83.58%) were male and 22 (16.42%) were female.

In present study wound infection [4 (16.66%)] was the most common complication followed by septicemia [2 (8.33%)].

**Table 1:** Distribution of patients according to incidence of various factors of BAT

Factors		2012 n=54 (%)	2013 n=50 (%)	2014 n=30 (%)
Age	0-10	5 (9.26)	6 (12)	0 (0)
	11-20	12 (22.22)	9 (18)	7 (23.33)
	21-30	13 (24.07)	14 (28)	7 (23.33)
	31-40	11 (20.37)	11 (22)	10 (33.33)
	41-50	2 (3.70)	7 (14)	5 (16.66)
	51-60	9 (16.66)	2 (4)	0 (0)
	>60	2 (3.70)	1 (2)	1 (3.33)
Treatment	Conservative	32 (59.29)	23 (43)	12 (40)
	Operative	22 (40.71)	27 (54)	18 (60)
EAI	Head Injury	15 (27.77)	8 (16)	8 (26.66)
	Chest	16 (29.62)	13 (26)	3 (9.99)
	Facial	1 (1.85)	8 (16)	0 (0)
	Pelvis	4 (7.40)	2 (4)	1 (3.33)
	Extremity	1 (1.85)	3 (6)	0 (0)
CP	Shock	4 (7.40)	12 (24)	2 (6.66)
	P/A Guarding tenderness	16 (29.62)	20 (40)	10 (33.33)
	Respiratory distress	14 (25.92)	10 (20)	5 (16.66)
	Localized tenderness	18 (33.33)	15 (30)	10 (33.33)
	Neurological sign	18 (33.33)	6 (12)	7 (23.33)
IOI	Liver	4 (7.40)	12 (24)	3 (9.99)
	Small intestine	6 (11.10)	15 (30)	6 (20)
	Spleen	4 (7.40)	9 (18)	4 (13.33)
	Hemoperitoneum	15 (27.77)	23 (46)	17 (56.61)
	KUB	2 (3.70)	4 (8)	3 (9.99)
	Other*	0 (0)	3 (6)	1 (3.33)

Data is expressed as no of patients (%), MOT; mode of trauma, RTA; road traffic accidents EAI; extra abdominal injuries, CP; clinical presentation, IOI; internal organ injuries, KUB; kidney, ureter and bladder, FBO; falling blunt object over body, \*stomach, pancreas and colon




Fig 1: Mode of trauma during study period in present study

Table 2: Distribution of patients according to other parameters

Parameters		2012 n=54 (%)	2013 n=50 (%)	2014 n=30 (%)
Hospital Stay (Days)	0-3	17 (31.48)	10 (20)	11 (36.67)
	4-6	12 (22.22)	11 (22)	3 (9.99)
	7-12	20 (37.03)	23(46)	15 (49.95)
	>12	5 (9.26)	6 (12)	1 (3.33)
LP	Liver Repair	1 (1.85)	3(6)	2 (6.66)
	ROP	6 (11.10)	10(20)	4 (13.32)
	RA	2 (3.70)	3 (6)	1 (3.33)
	Splenectomy	2 (3.70)	0 (0)	0 (0)
	EH	8 (14.8)	7 (14)	5 (16.65)
	KR & SPC	2 (3.70)	1 (2)	2 (6.66)
Complication	Other*	0 (0)	4 (8)	1 (3.33)
	WI	0 (0)	2(4)	2 (6.66)
	Septicemia	0 (0)	4 (8)	0 (0)
	Burst abdomen	0 (0)	1 (2)	0 (0)
	Other^	0 (0)	0 (0)	0 (0)
Outcome	Death	6 (11.10)	13 (26)	5 (16.65)
	IWOC	48 (88.89%)	33 (66)	23 (76.59)
	IWC	0 (0)	4 (8)	2 (6.66)

Data is expressed as no of patients (%), ROP; repair of perforation, RA; resection & anastomosis, EH; evacuation of hemo peritoneum; KR & SPC: Kidney Repair & Supra Pubic catheterization PC, WI; wound infection, IWOC; Improved without complication, IWC; improved with complication; M & M; mortality and morbidity, LP; laprotomy procedure\*mesenteric/diaphragm tear repair/ileostomy/colostomy;^ DVT, Pulm Complication, Hepato/renal Complication, bed sore

**Discussion**

BAT is still one of the major reasons of trauma admission throughout world. It also significantly leads to high morbidity and mortality [9].

In present study there was a decline in the incidence of BAT from 2012 to 2014 which may be due to combination of factors including increase in effective road safety policies and decrease in violence in urban area which is consistency with the study done by Asuquo *et al.* [10].

In harmony with other studies done by Manohar *et al.*, Asuquo *et al.*, Kulkarni *et al.*, and Chalya *et al.*, most of the patients with BAT in present study were young in their second and third decade of life and affected mostly men as compared to female [9, 10, 11, 12].

Patients having age between 20-30 year represents the economically active group and this is the reason why there is a high incidence of BAT in this age group. The reason behind the male predominance in present study is due high

engagement of men in high-risk activities. In most of the Indian household male population are bread earners and likely to involve in such activities which can lead to injury in pursue of attempting to earn a living.

Asuquo *et al.*, performed a similar study on 19 BAT patients and reported road traffic accidents as the most common cause for BAT which is consistency with the findings of present study. [10, 13] Study done by Kulkarni *et al.*, including 68 consecutive cases of blunt trauma and another study done by Jolly *et al.*, also reported road traffic accident as the most common cause of BAT [11, 14].

Srihari *et al.*, performed a similar study and revealed that among all BAT patients most of them were managed surgically whereas only 31.6% were treated conservatively. Similarly in present study most of the patients were treated conservatively in 2012 but trend was shifted to surgical intervention during 2013 and 2014 which is similar to the reports by Srihari *et al.* [6].

Head injury and chest were the most common extra abdominal injuries in present study and most of the patients were presented with localized tenderness which is consistency with the study performed by Chalya *et al.* [12] Damage to the visceral organ or fatal hemorrhage in BAT patients makes it one of the important types of trauma which require correct diagnosis along with early intervention. Doing so may reduce the chances of death and associated morbidity [11, 15]. In present study small intestine was the most commonly involved internal organ which is consistent with the study done by Kulkarni S 2015 [11].

Srihari *et al.*, reported 13.33% mortality septicemia being the most common cause of death in their study. Similar data was reported in present study [6]. In present study most common complication was wound infection followed by septicemia. Similar results were depicted by Naveen *et al.*, and Srihari *et al.* [6, 16].

Laparotomy was performed in patients to locate and repair injured organ, to clean peritoneal cavity and to look over for any other abdominal cavity injuries. In present study hemoperitoneum was the most commonly observed intra-operative finding, similar results were reported by Chalya *et al.* [12] Present study has few limitation of being small in size; a large randomized clinical trial is required to confirm the results.

**Conclusion**

Average incidence of BAT in present study was 0.72% with male predominance. Most of the patients were in young age. Head injury and chest were the most common extra abdominal injuries, small intestine was mainly got damaged, most of the patients stated for 7 to 12 days in the hospital and though many patients died due to severity of the trauma, most of them recovered without any complications.

**References**

- Poletti PA, Kinkel K, Vermeulen B, *et al.* Blunt abdominal trauma: should US be used to detect both free fluid and organ injuries? *Radiology.* 2003; 227:95-103.
- Livingston DH, Lavery RF, Passannante MR, *et al.* Admission or observation is not necessary after a negative abdominal computed tomographic scan in patients with

- suspected blunt abdominal trauma: results of a prospective, multi-institutional trial. *J Trauma* 1998; 44:273-280.
3. Kendall JL, Kestler AM, Whitaker KT, Adkisson MM, Haukoos JS. Blunt Abdominal Trauma Patients Are at Very Low Risk for Intra-Abdominal Injury after Emergency Department Observation. *Western Journal of Emergency Medicine* 2011; 12(4):496-504.
  4. Poletti PA, Mirvis SE, Shanmuganathan K, *et al.* Blunt abdominal trauma patients: can organ injury be excluded without performing computed tomography? *J Trauma*. 2004; 57:1072-1081.
  5. Stephan PJ, McCarley MC, O'Keefe GE, *et al.* 23-Hour observation solely for identification of missed injuries after trauma: is it justified? *J Trauma*. 2002; 53:895-900.
  6. Srihari V, Jayaram J, Sabira S. A Clinical Study of Blunt Injury Abdomen. *Indian Journal of Research*. 2015; 4(1):123-6.
  7. Pimentel SK, Sawczyn GV, Mazepa MM, Gonçalves Da Rosa FG, Nars A, Collaço IA. Risk factors for mortality in blunt abdominal trauma with surgical approach. *Rev Col Bras Cir*. 2015; 42(4):259-64.
  8. Cowell VL, Ciraulo D, Gabram S, *et al.* Trauma 24-hour observation critical path. *J Trauma*. 1998; 45:147-50.
  9. Manohar K, Ramanaiah GV. Abdominal Trauma in adults - its outcome-a prospective study in a tertiary health care centres in Andhra Pradesh. *Indian Journal of Applied Research*. 2015; 5(11):35-8.
  10. Asuquo M, Nwagbara V, Umoh M, Ugare G, Agbor C, Japhet E, *et al.* Blunt Abdominal Trauma in a Teaching Hospital, Calabar, Nigeria. *International Journal of Clinical Medicine*. 2012; 3:693-6.
  11. Kulkarni S, Kanase V, Kanase N, Varute P. Blunt Trauma to Abdomen in Rural Setup: A Multiple Case Study. *Int J Sci Stud*. 2015; 3(4):16-9.
  12. Chalya PL, Mabula JB. Abdominal trauma experience over a two-year period at a tertiary hospital in northwestern Tanzania: a prospective review of 396 cases. *Tanzania Journal of Health Research*. 2013; 15 (4):1-13.
  13. Dogo D, Yawe T, Hassan AW, Tahir B. Pattern of Abdominal Trauma in North Eastern Nigeria. *Nigerian Journal of Surgical Research*. 2000; 2:48-51.
  14. Jolly S, Upadhyay M, Jain BL. Blunt abdominal trauma. A clinical study of 100 cases. *Indian J Surg*. 1993; 6:55.
  15. Kennedy RH. Non penetrating injuries of abdomen. *AMA Arch Surg*. 1957; 75:957-63.
  16. Naveen KG, Ravi N, Nagaraj B. Blunt Abdominal Trauma: Making Decision of Management with Conventional and Ultrasonography Evaluation. *SSRG International Journal of Medical Science (SSRG-IJMS)*. 2014; 1(2):1-13.