



Study of factors related to time of breastfeeding initiation among mother-infant pairs in Ibn al-Baladi hospital in Baghdad Iraq

Dr. Jenan Ghadban Dawood¹, Dr. Talib Hasan Darhash Al Tamimi², Dr. Kaddum Kudair Abbas AL Taie³

¹ MBChB, DCH, College of Medicine, AL Mustansiriya University Baghdad, Iraq

² MBChB, DCH, Kufa College of Medicine, Baghdad University, Iraq

³ Department of Pediatrics-Ibn Baladi Hospital- Baghdad, Iraq

Abstract

Early initiation of breastfeeding (EIBF) and exclusive breastfeeding for the first 6 months of life prevents around 20% newborn deaths and 13% under-five deaths. EIBF is predominately associated with socioeconomic, health related and individual factors. This study was carried out with the objective to find out the prevalence of EIBF and association of demographic, antenatal and intra natal factors with pre lacteal feeding practice in Al-Sadir city in Baghdad.

This study was carried out in 2017–2018 on a sample of 200 mother-infant pairs calculated with appropriate formula using EIBF prevalence. Multistage sampling was done and mothers who gave valid consent were taken as respondent. Chi-square, Fisher exact and Pearson correlation tests were used for statistical analysis.

The prevalence of EIBF was 65% in Al-Sadir city. About 93.1% newborn of 15–19 years aged mothers, 96.4% newborns of mothers of Class-V socio-economic class, 87.5% newborns of illiterate mothers, 93.8% newborns of farmer mothers, and 78.9% of 1st birth order newborns received breastfeeding within 1 h of their birth. The proportion of newborn receiving breastfeeding within 1–4 h after birth increased significantly with increasing number of antenatal care (ANC) visits ($P < 0.05$). Among mothers with prolonged labor, 43.5% could initiate breastfeeding early. Mode of delivery and health facility for delivery were a significant determinant for the time of initiation of breastfeeding.

EIBF decreased with increasing age of mothers, increasing socio-economic class and increasing the education level of mothers. The practice of EIBF is also low among employed mothers, among newborn with high birth order and among mothers of general caste. EIBF is high in mothers with fewer ANC visits, among mothers with prolonged labor, in case of normal delivery, delivery at primary health care, and delivery assisted by female health staff.

Keywords: early initiation; breastfeeding; socio-demographic factors; antenatal care

1. Introduction

Every infant and child has the right to good nutrition according to the “Convention on the Rights of the Child.” Undernutrition is associated with 45% of child deaths [1]. Globally in 2016, 155 million children under 5 years of age were estimated to be stunted, 52 million were estimated to be wasted, and 41 million were overweight or obese [1].

Infant feeding practices are one of the determinants for the nutritional status of a child [2]. Recent scientific evidence reveals that malnutrition has been responsible directly or indirectly for 60% of all deaths among under-five age children annually. Inappropriate feeding practices results in over two-third of under-five deaths. Poor feeding practice in infancy and childhood, resulting in malnutrition contribute to impaired cognitive and poor social development, poor school performance, and reduced productivity in later life. WHO and United Nations Children’s Fund (UNICEF) recommend early initiation of breastfeeding (EIBF) within 1 h of birth [3], exclusive breastfeeding for the first 6 months of life; and introduction of nutritionally-adequate and safe complementary foods at 6 months together with continued breastfeeding up to 2 years of age or beyond [1]. EIBF and exclusive breastfeeding for the first 6 months of life prevent around 20% newborn deaths and 13% under-five deaths [4]. It can also reduce mortality due to neonatal infections (sepsis, pneumonia, tetanus, and diarrhea) [5], which contribute to

36% in neonatal deaths from all causes [6]. The first breast milk is known as colostrum, which is highly nutritious and contains antibodies that protect the newborn from diseases [3]. Many newborns are given prelacteal feeding before starting breastfeeding. It consists of honey, jaggery, sugar water, etc. This may be specific to a family, caste, or religion [7]. The harmful effects of prelacteal feeding include a delay in the initiation of breastfeeding and prevention of the initial bonding between the mother and the baby [8]. With discarding colostrum and delaying breastfeeding, a marked increase in neonatal mortality has been reported from different parts of the world [9]. The deleterious effects of infections causing infant deaths can be prevented by EIBF and exclusive breastfeeding, which is the easiest, cost-effective, and life-saving intervention for the health of a newborn [10].

Worldwide countries are scaling up efforts to increase rates of optimal breastfeeding practices to decrease child mortality. Breastfeeding is linked to many objectives of the 2030 agenda for Sustainable Development Goals. World Breastfeeding Week is celebrated every year for raising awareness regarding the fact that breastfeeding is a key to sustainable development. India has made great strides in improving child nutrition over the past decade. The government of India has promoted child health programs and is advocating healthy infant child feeding practices. One such program is the implementation of the “National Infant and

Young Child Feeding” Guideline, which discourages prelacteal feeding practices and encourages optimal breastfeeding practices. The concept of baby friendly hospital initiative (BFHI) was launched globally in 1991 by UNICEF and the WHO for promoting and supporting breastfeeding. Including India, many hospitals from different countries have been designated to BFHI. In the year of 2016, the Government of India launched the National Breastfeeding Promotion Programme mothers’ absolute affection to ensure adequate awareness is generated among the masses, especially mothers, on the benefits of breastfeeding. The programme will be implemented at three levels: Macrolevel through mass media; meso level in health facilities, and microlevel at communities [11].

Prior studies have shown that EIBF is associated with a lower risk of neonatal mortality [12-14]. It is found globally that only half of newborn babies are breastfed during their 1st h of birth, despite strong evidence of nutritional and immunological benefits of early initiation in reducing neonatal mortality and morbidity [14-16]. The first milk (colostrum) contains bioactive immune factors which protect a neonate against a variety of infections and allergic diseases [17]. A recent systematic review of literature based on 25 studies from seven countries in South Asia revealed that EIBF is predominately associated with socioeconomic, health related, and individual factors [18].

We have carried out this study with the objective to know the role of women’s demographic factors; antenatal care (ANC) and delivery care with the timing of initiation of breastfeeding to their newborn.

Materials and methods

This community-based cross-sectional retrospective study was carried out during the period from September 2017 to August 2018. Mothers-infant pairs who were delivered during the past 6 months before the period of data collection were the respondents.

Household visits of mothers were carried out with prior telephonic intimation. In case of locked house or nonresponse, we excluded that mother from the study. After taking the informed consent, detailed interview of mother was conducted, and answers were recorded in the pre-designed pre-validated structured questionnaire. Feeding practice of the latest born child was asked.

Study tool consisted of structured questionnaire with two parts. First part recorded the socio-demographic information including age, education and occupation of mothers, type and size of family, caste, family income, questions for socioeconomic class, etc. The second part of the questionnaire consisted of timing of initiation of breastfeeding and ANC. The timing of initiation of breastfeeding was defined as when after birth, the mother-initiated breastfeeding. We used the term EIBF to indicate initiation within 1 h of delivery according to the WHO recommendation and late initiation to indicate after 1 h of birth [19].

The following criteria were included in the study: Mothers who delivered during the past 6 months at the time of interview, mothers whose child was currently alive and mothers who gave valid consent. While mothers who did not give valid consent were excluded from the study.

Statistical Methods

Data were analyzed using Microsoft Excel and SPSS v20. Chi square tests (χ^2) and Fisher exact tests were performed to evaluate the association of the independent variables with the EIBF. Pearson correlation test was used to examine linear correlation for quantitative variables.

Results

The mean and standard deviation of mother’s age is 24.95 ± 6.31 years. Distribution of newborns according to time of initiation of breastfeeding after birth shows that 65% newborn received breastfeeding within 1 h of birth, i.e., EIBF. The proportion of newborns who received breastfeeding between 1 and 4 h after birth and more than 4 h after birth was 20% and 15%, respectively [Figure 1].




Fig 1

Analysis of relation of demographic characteristics of mother with time of initiation of breastfeeding after birth shows that highest 93.1% mothers in age group of 15–19 years-initiated breastfeeding within 1 h of birth as compared to lowest 20.0% mothers in age group of 35–39 years which is also statistically significant ($P < 0.005$). A statistically significant difference in EIBF was observed across different socio-economic classes ($P < 0.05$). EIBF was highest (96.4%) among mothers of Class-V family followed by Class-II with 70.2%. Education of mother wise analysis shows that EIBF was highest (87.5%) among illiterate while no one from graduate mothers could initiate it early ($P < 0.01$). Occupation wise only 30.4% of employed mothers could initiate early breastfeeding. Interestingly 93.8% of farmers and 73.2% housewife initiated breastfeeding within 1 h of birth ($P < 0.005$). Statistically significant decrease in the prevalence of EIBF practice was observed with increased parity ($P < 0.005$). The highest prevalence of EIBF was found among those mothers having one child and lowest in mothers with previous four children [Table 1].

It is evident from Table 2 that 77.8% of mothers who were neither registered during the antenatal period and nor received ANC had initiated breastfeeding within 1 h of birth. However, 20.9% of those mothers who were registered and received ANC initiated breastfeeding within 1–4 h duration after delivery as compared to only 11.1% of mothers who were not registered and not received ANC. The difference in a time of initiation of breastfeeding among ANC received mothers, and those not received ANC is not significant

Statistically (P=0.111). Interestingly 94.1% of mothers who received only one ANC initiated breastfeeding within 1 h as compared to lowest 47.4% mothers who received ANC 5 times. Analysis of source of ANC shows 80% mothers who received ANC from Community Health Centre (CHC)/district hospital initiated within 1 h of delivery as compared to 61% of those mothers who received ANC from PHC/Urban Health Centre. Analysis of the impact of complication on time of initiation of breastfeeding shows that 43.5% of mothers who had prolonged labor could practice EIBF. There is statistically significant (P=0.005) difference in time of initiation breastfeeding among mother without complications and those with other complications such as

bleeding and hypertension. It is observed that 75.5% of those mothers who had normal delivery initiated breastfeeding within 1 h as compared to 12.2% of those mothers who had delivery by cesarean section (P=0.005). There is no statistically significant difference in the time of initiation of breastfeeding between institutional and home delivery (P=0.738). However, within institutional deliveries, the difference is statistically significant (P=0.005). Among mothers, who delivered baby at PHC; 96.8% of them initiated breast feeding early while 44.4% of mothers who delivered at private hospital initiated early breast feeding. The assistance of delivery by Auxiliary Nurse-Midwife (ANM) or staff nurse led to EIBF among 76.0% newborns.

Table 1: Relation of initiation of breastfeeding with demographic details of pregnant women (n=200)

	Breastfeeding initiated (h)			Total No. (%)	Statistical results
	<1 No. (%)	1-4 No. (%)	>4 No. (%)		
Age group					
15-19	27(93.1)	0 (0)	2 (6.9)	29	$\chi^2=131.65, P<0.005$
20-24	40(64.5)	22 (35.5)	0 (0)	62	
25-29	45(73.8)	16 (26.2)	0 (0)	61	
30-34	10(26.3)	3 (7.9)	25(65.8)	38	
35-39	2(20.0)	0 (0)	8(80.0)	10	
Socio -economic class					
Class-1	6(20.0)	1(3.3)	23(76.7)	30	$\chi^2=102.46, P<0.005$
Class-11	40(70.2)	11(19.3)	6(10.5)	57	
Class-111	28(54.9)	18(35.3)	5(9.8)	51	
Class-1v	23(67.6)	10(29.4)	1(3.0)	34	
Class-v	27(96.4)	1(3.6)	0(0)	28	
Education of mother up to					
Graduate	0 (0)	0 (0)	18(100)	18	$\chi^2=153.22, P<0.005$
Higher secondary	7 (26.9)	3(11.5)	16(61.6)	26	
Secondary	40 (69.0)	18(31.0)	0 (0)	58	
Primary	28 (82.4)	4(11.8)	2(5.8)	34	
Just illiterate	28 (70.0)	12(30.0)	0 (0)	40	
Illiterate	21 (87.5)	3(12.5)	0 (0)	24	
Occupation					
Employed	14 (30.4)	4(8.7)	28(60.9)	46	$\chi^2=107.97, P<0.005$
Farmer	30 (93.8)	0 (0)	2(6.2)	32	
Housewife	60 (73.2)	20(24.4)	2(2.4)	82	
Labourer	14 (53.8)	12(46.2)	0 (0)	26	
Student	3 (21.4)	7(50.0)	4(28.6)	14	
Birth order					
1st	45 (78.9)	10(17.5)	2(3.6)	57	$\chi^2=125.35, P<0.005$
2nd	59 (73.8)	21(26.2)	0 (0)	80	
3rd	18 (45.0)	9(22.5)	13(32.5)	40	
4th	1 (6.7)	0 (0)	14(93.3)	15	
5th	0 (0)	0 (0)	8(100)	8	

Table 2: Relation of initiation of breastfeeding with ANC and obstetric care of pregnant women (n=200)

ANC/Obstetric care	Breastfeeding initiated (h)			Total	Statistical results
	<1	1-4	>4		
ANC/registered					
Yes	117(61.3)	40 (20.9)	34 (17.8)	191	$\chi^2=1.0, P=0.625$
No	7(77.8)	1 (11.1)	1 (11.1)	9	
Number of times ANC received					
0	7 (77.8)	1(11.1)	1(11.1)	9	$\chi^2=15.62, P=0.111$
1	16 (94.1)	1 (5.9)	0 (0)	17	
2	12 (80.0)	3 (20.0)	0 (0)	15	
3	50 (56.8)	19 (21.6)	19 (21.6)	88	
4	30 (57.7)	12 (23.1)	10 (19.2)	52	
5	9 (47.4)	5 (26.3)	5 (26.3)	19	
Source of ANC					
PHC/UHC	64 (61.0)	22 (21.0)	19 (18.0)	105	$\chi^2=1.52, P=0.823$
Sub center/	59 (65.6)	17 (18.9)	14 (15.5)	90	
CHC/District hospital	4 (80)	1 (20)	0 (0)	5	

Pregnancy-related complications					
No complication	82 (60.3)	20 (14.7)	34 (25.0)	136	$\chi^2=32.45, P=0.005$
Bleeding	13 (68.4)	6 (31.6)	0 (0)	19	
Convulsion	6 (60.0)	4 (40.0)	0 (0)	10	
Hypertension	12 (100)	0 (0)	0 (0)	12	
Prolonged labor	10 (43.5)	11 (47.8)	2 (8.7)	23	
Type of delivery					
Normal	120 (75.5)	38 (23.9)	1 (0.6)	159	$\chi^2=152.93, P=0.005$
Cesarean section	5 (12.2)	2 (4.9)	34 (82.9)	41	
Place of delivery					
Institutional	118 (61.5)	40 (20.8)	34 (17.7)	192	$\chi^2=0.61, P=0.738$
Home	6 (75.0)	1 (12.5)	1 (12.5)	8	
Health facility utilized for delivery					
CHC	18 (58.1)	13 (41.9)	0 (0)	31	$\chi^2=47.08, P=0.005$
District hospital	52 (55.9)	17 (18.3)	24 (25.8)	93	
PHC	30 (96.8)	1 (3.2)	0 (0)	31	
Private hospital	20 (44.4)	10 (22.2)	15 (33.4)	45	
Assisted delivery					
ANM/Staff nurse	38 (76.0)	12 (24.0)	0 (0)	50	$\chi^2=15.97, P=0.013$
TBA	1 (33.3)	1 (33.3)	1 (33.3)	3	
Doctor	80 (55.9)	28 (19.6)	35 (24.5)	143	
Untrained	2 (80)	1 (10)	1 (10)	4	

UHC: Urban health center, PHC: Primary health care, TBA: Trained birth attendant

Discussion

The prevalence of EIBF was 65%. About 93.1% newborn of 15–19 years aged mothers, 96.4% newborns of mothers of Class-V socio-economic class, 87.5% newborns of illiterate mothers, 93.8% newborns of farmer mothers, and 78.9% of 1st birth order newborns received breastfeeding within 1 h of their birth. There is no statistically significant difference in timing of initiation of breastfeeding among mothers who received antenatal and who did not receive it. However, with the increasing number of ANC visits, the proportion of newborn receiving breastfeeding within 1–4 h after birth increased. Among mothers with prolonged labor, 43.5% could initiate breastfeeding early. Mode of delivery and health facility for delivery were a significant determinant for the time of initiation of breastfeeding.

NFHS-4 conducted in 2015–2016 has shown the total prevalence of EIBF as 49.9% in Gujarat [21] and 52.1% in a rural area of Jamnagar district [21]. Gandhi *et al.* in their study conducted in Navsari district of Gujarat in 2012–13 found that 56.4% of mothers initiated breastfeeding within 1 h. [20]. Early initiation of exclusive breastfeeding serves as the starting point for a continuum of care for mother and newborn that can have long-lasting effects on health and development.[26] We found a significantly higher prevalence (93.1%) of EIBF practice in mothers with 15–19 years age group than any other age groups ($P < 0.05$). Similarly Islam *et al.* in their nationwide study conducted in 2014 in Bangladesh also found that the rate of EIBF was significantly ($P < 0.05$) higher (52.5%) among those who were married before 18 years as compared to those mothers who were married after 18 years of age (48.5%) [27]. They also found that there was a significantly ($P < 0.05$) higher rate of EIBF in newborns of mothers with <20 years of age (53.0%) when compared to those who were older than 20 years (46.9%) when they delivered their first child [27]. Garje *et al.* in Kerala found that 74.3% mothers in the age group of 21–25 years initiated breastfeeding within 30 min as compared to only 6.6% of those mothers having more than 26 years of age [23]. Younger mothers are more receptive to the advice given by health worker as well as they have less psychological stress, which may lead to EIBF. Physiologically, breast milk

secretion decreases with increasing age. Dewey *et al.* reported that lactational capacity is greatest before age 20, and declines thereafter [28]. They further stated that “older” primiparae (24 years or older) produced less milk in the 1st week postpartum than younger primiparae (16–23 years) [28]. Garje *et al.* demonstrated that 53.6% of schedule caste mothers initiated breastfeeding within 30 min and 69.1% within 30 min to 1 h as compared to 4.9% and 3.1% of general caste, respectively [23]. Gandhi *et al.* also found similar results with 67.2% prevalence of EIBF among scheduled tribe mothers. Increasing prevalence of EIBF was observed as moving down from higher to lower socio-economic classes. Similarly, Islam *et al.* in Bangladesh observed that 54.9% of families with poor wealth index practiced EIBF as compared to 47% of those with rich wealth index [27]. Mothers of higher socio-economic class may be more stressful, which may lead to delay in the secretion of breast milk. Moreover, we found that higher education among mothers was not a favoring factor for EIBF. The proportion of EIBF practice was higher among illiterate and just illiterate mother as compared to those who were graduate or had studied up to higher secondary class ($P < 0.05$). Similarly, Islam *et al.* also reported that education had a negative influence on EIBF, implying that mothers with secondary education and higher education were less likely to practice EIBF than those who were uneducated ($P < 0.01$) [27]. Garje *et al.* found that 44.8% and 41% mothers with primary and secondary education, respectively, initiated breastfeeding within 30 min of delivery as compared to 9.3% and 1.1% of those with who were illiterate and undergraduate, respectively [23]. Higher education women are mostly employed and employment stress may delay secretion of breast milk secretion. Moreover, highly educated and employed women may be more conscious about the cosmetic outlook and may have myths that breastfeeding may disfigure them. Women with lower education, housewives, farmers, and laborers may be more receptive to feeding advice given by health-care provider. That is why 93.8% of farmer mothers, 73.2% of housewives practiced EIBF as compared to 30.4% of employed mothers. Similarly, Garje *et al.* reported that 59.6% of daily wage women, 34.4% of farmers and 6% of business

women initiated breastfeeding within 30 min of delivery [23]. Various other studies conducted in various parts of India support our findings of relations of socio-demographic factors with EIBF [24, 28-30]. Higher proportion of EIBF was found among newborns with birth order 1st and 2nd as compared with those with birth order three or more ($P < 0.05$). In contrast to that Gandhi *et al.* in Navsari district of Gujarat found 48.5% of 1st birth order newborn and 65% of 2nd and 3rd birth order newborns were given early breastfeeding as compared to 83.5% of those with birth order four and above [20]. We presume that the age of mother would be also more for those newborn with high birth order in our study and increased age has a negative effect on EIBF in our study. High birth order also impairs the nutritional status of mother leading to decreased and delayed breast milk secretion.

ANC is not statistically significantly associated with EIBF. One of the reasons for this may be small sample size (i.e., 9) in the category of those mothers who did not receive ANC, and second, they may have received advice for EIBF from relatives, neighbors and even during delivery. ANC services basically focus on maternal health. Islam *et al.* also found that mothers who regularly attended the antenatal clinic (3 and more times) were 1.24 times more likely to practice EIBF compared to mothers who did not attend the ANC visits ($P < 0.05$). An increasing number of antenatal visits give more opportunity to health-care provider to counsel mother about newborn care in addition to maternal care. Naturally, intra natal complications such as prolonged labor delay initiation of breastfeeding to newborn as found in our study; however, antenatal complications such as bleeding, convulsion, and hypertension individually were not inhibiting factors for EIBF. Islam *et al.* found that 58.3% of those mothers with normal delivery practiced EIBF as compared to 30.1% of those delivered by cesarean section [27]. Our findings are also in line with this study. It may take time for recovery from anesthesia and trauma in case of cesarean delivery, which delays breastfeeding. Place of delivery although not significantly associated for EIBF; our finding of a higher proportion of EIBF (75%) in case of home delivery as compared to institutional delivery (61.5%) is supported by the study of Islam *et al.* who found 58.8% EIBF in home delivery versus 40.6% in institutional delivery [27]. Mother may feel more privacy at home for feeding of newborn. Health facility where mother delivered was found to be associated with EIBF in our study; as high as 96.8% of those mothers who were delivered at PHC practiced EIBF ($P < 0.05$). Mothers who deliver in PHC have normal vaginal delivery while complicated cases are generally referred to CHC or higher level where due to cesarean section, breastfeeding may be delayed. Assistance of delivery by TBA and ANM was a favoring factor for EIBF. TBA are also now aware of the benefits of EIBF and ANM being female gender can easily communicate with the mother for feeding advice.

References

1. World Health Organization. Infant and Young Child Feeding Key Facts. Geneva: World Health Organization; 2018.
2. Betrán AP, de Onís M, Lauer JA, Villar J. Ecological study of effect of breast feeding on infant mortality in Latin America. *BMJ*. 2001; 323:303-6.
3. Acharya R, Meena R. A descriptive cross-sectional study of breast-feeding practice in Bikaner, Rajasthan. *Int J Med Sci Public Health*. 2016; 5:1559.
4. Jones G, Steke tee RW, Black RE, Bhutta ZA, Morris SS. Bellagio Child Survival Study Group, *et al.* How many child deaths can we prevent this year? *Lancet*. 2003; 362:65-71.
5. Mull any LC, Katz J, Li YM, Khattry SK, Le Clerq SC, Darmstadt GL, *et al.* Breast-feeding patterns, time to initiation, and mortality risk among newborns in Southern Nepal. *J Nutr*. 2008; 138:599-603.
6. Lawn J, Shibuya K, Stein C. No cry at birth: Global estimates of intra partum stillbirths and intra partum-related neonatal deaths. *Bull World Health Organ*. 2005; 83:409-17.
7. Shankar K, Muthu kumar S. Pre lacteal feeding practice among rural mothers in Tamilnadu. *Int J Biomed Adv Res*. 2015; 6:484-7.
8. Parashar A, Sharma D, Gupta A, Dhadwal DS. Associated factors in Himachal Pradesh. *Int J Health*. 2017; 6:30-4.
9. Jayarama S, Ramaiah R. Pre lacteal feeding practice among mothers in a rural area of Karnataka: A cross sectional study. *Int J Community Med Public Health*. 2017; 4:2919.
10. Lutter C. Early Initiation of Breastfeeding: The Key to Survival and Beyond. Healthy Life Course Project. Washington D.C: Pan American Health Organization, 2010.
11. MAA-Mothers' Absolute Affection. Operational Guideline. New Delhi: Ministry of Health and Family Welfare, 2016.
12. Berde AS, Yalcin SS. Determinants of early initiation of breastfeeding in Nigeria: A population-based study using the 2013 demographic and health survey data. *BMC Pregnancy Childbirth*. 2016; 16:32.
13. Debes AK, Kohli A, Walker N, Edmond K, Mullany LC. Time to initiation of breastfeeding and neonatal mortality and morbidity: A systematic review. *BMC Public Health*. 2013; 13(3):S19.
14. NEOVITA Study Group. Timing of initiation, patterns of breastfeeding, and infant survival: Prospective analysis of pooled data from three randomised trials. *Lancet Glob Health*. 2016; 4:e266-75.
15. Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J, *et al.* Breastfeeding in the 21st century: Epidemiology, mechanisms, and lifelong effect. *Lancet*. 2016; 387:475-90.
16. Khan J, Vesel L, Bahl R, Martines JC. Timing of breastfeeding initiation and exclusivity of breastfeeding during the first month of life: Effects on neonatal mortality and morbidity a systematic review and meta-analysis. *Matern Child Health J*. 2015; 19:468-79.
17. Chae A, Aitchison A, Day A, Keenan JI. Bovine colostrum demonstrates anti-inflammatory and antibacterial activity in *in vitro* models of intestinal inflammation and infection. *J Funct Foods*. 2017; 28:293-8.
18. Sharma IK, Byrne A. Early initiation of breastfeeding: A systematic literature review of factors and barriers in South Asia. *Int Breastfeed J*. 2016; 11:17.
19. Bhandari DJ, Pandya YP, Sharma DB. Barriers to exclusive breastfeeding in rural community of central Gujarat, India. *J Fam Med Prim Care*. 2019; 8:54-61.
20. Gandhi S, Godara N, Modi A, Kantharia S. Impact of feeding practices on nutritional status of children in rural

- area of Navsari district. *Int J Med Sci Public Health*. 2014; 3:1338.
21. International Institute for Population Sciences. National Family Health Survey-4. India, Mumbai: International Institute for Population Sciences, 2017.
 22. World Health Organization (WHO). Indicators for Assessing Infant and Young Child Feeding Practices: Conclusions of a Consensus Meeting Held 6-8 November 2007 in Washington D.C., USA: World Health Organization (WHO), 2008.
 23. Garje R, Thitame S, Somasundaram K, Gautam L, Gite M. Infant feeding beliefs and practices among tribal mothers in Nashik district, Maharashtra. *Int J Med Sci Public Health*. 2016; 5:2203.
 24. Manghat S, Thulaseedharan J. Feeding practices during first 6 months of infancy observations from a community-based cross-sectional survey in central Kerala, India. *Int J Med Sci Public Health*. 2018; 7:434-9.
 25. Gupta A, Dadhich JP, Ali SM, Thakur N. Skilled counseling in enhancing early and exclusive breastfeeding rates: An experimental study in an urban population in India. *Indian Pediatr*. 2019; 56:114-8.
 26. Insight. Impact of Early Initiation of Exclusive Breastfeeding on Newborn Deaths. Washington, DC: A and T Technical Brief, 2010.
 27. Islam MA, Mamun A, Hossain MM, Bharati P, Saw A, Lestrel PE, *et al*. Prevalence and factors associated with early initiation of breastfeeding among Bangladeshi mothers: A nationwide cross-sectional study. *PLoS One*. 2019; 14:e0215733.
 28. Dewey K, Finley D, Strode M, Lönnerdal B. Relationship of Maternal Age to Breast Milk Volume and Composition. *Human Lactation*. Boston, MA: Springer Link. 1986; 2:263-73.
 29. Singh S, Tiwari N, Malhotra A. A cross-sectional study on exclusive breastfeeding practice among lactating females attending medical college, district Jhansi (U.P). *Int J Med Sci Public Health*. 2017; 6:377-8.
 30. Bashir A, Mansoor S, Naikoo M. Knowledge, attitude, and practices of postnatal mothers regarding breastfeeding: A cross-sectional study. *Int J Med Sci Public Health*. 2018; 7:725.